

Degeneracy in Density Functional Theory: Topology in v - and n -Space

C. A. Ullrich* and W. Kohn

Department of Physics, University of California, Santa Barbara, CA 93106

(Dated: February 1, 2008)

This paper clarifies the topology of the mapping between v - and n -space in fermionic systems. Density manifolds corresponding to degeneracies $g = 1$ and $g > 1$ are shown to have the same mathematical measure: every density near a g -ensemble- v -representable (g -VR) $n(\mathbf{r})$ is also g -VR (except “boundary densities” of lower measure). The role of symmetry and the connection between $T = 0$ and $T = 0^+$ are discussed. A lattice model and the Be-series are used as illustrations.

PACS numbers: 31.15.Ew, 31.15.Ar, 71.10.Ca, 71.15.Mb

Density functional theory (DFT) of electrons in an external potential $v(\mathbf{r})$ uses the ground-state density $n(\mathbf{r})$ as basic variable [1, 2]. $n(\mathbf{r})$ uniquely determines $v(\mathbf{r})$ (apart from a constant); conversely, for a given number, N , of electrons, $v(\mathbf{r})$ obviously determines $n(\mathbf{r})$ uniquely if and only if the ground state is non-degenerate. These facts point to the interesting role of degeneracy in the mapping $v(\mathbf{r}) \longleftrightarrow n(\mathbf{r})$.

In 1982-3, Levy [3] and Lieb [4] discovered examples of well-behaved density functions, $n(\mathbf{r})$, which *could not* be reproduced as non-degenerate, non-interacting ground-state densities of *any* $v(\mathbf{r})$, and were called non- v -representable (non-VR). In these examples, the $n(\mathbf{r})$ *could* be reproduced as weighted averages of densities of degenerate ground states corresponding to a $v(\mathbf{r})$.

Soon afterwards, Kohn [5] adopted a lattice version of the Schrödinger equation, $\mathbf{r} \rightarrow r_l$ ($l = 1, \dots, M$), in which $v(\mathbf{r})$ and $n(\mathbf{r})$ become v_l and n_l and can be viewed as M' -dimensional (M' -D) vectors: $\vec{v} \equiv (v_1, \dots, v_{M'})$, with $M' \equiv M - 1$, $v_M = 0$; and $\vec{n} \equiv (n_1, \dots, n_{M'})$, with $n_M = N - (n_1 + \dots + n_{M'})$ [6]. The M lattice points are enclosed by boundary points on which $\vec{v} = +\infty$ and, accordingly, all wave functions vanish. It was shown in [5] that in the M' -D $\vec{n}$ -space there are finite M' -D regions in which all “points” (densities) are VR.

In 1985, Chayes *et al.* (CCR) [7], using a similar lattice model, proved the following important result: Any well-behaved $\vec{n}$ ($0 < n_l \leq 1$) can be uniquely represented as weighted average of g degenerate ground-state densities associated with a $\vec{v}$. We call the special case $g = 1$ 1-VR, and the general case ($g \geq 1$) g -VR. Thus, $\vec{n}$ -space is filled by manifolds S^g , in each of which $\vec{n}$ is g -VR. Each S^g maps on a corresponding manifold Q^g in $\vec{v}$ -space.

This paper aims at an understanding of the topologies of the regions S^g and Q^g and of the nature of the boundaries between them. This has mathematical and physical significance, e.g. in the search for self-consistent solutions of Kohn-Sham (KS) equations [8]. We shall see that g -VR densities with $g > 1$ are *not* mathematically exceptional and, in particular, do *not* depend on symmetries of the potential. As illustrations we shall present a finite lattice model and the Be-series for $Z \rightarrow \infty$ [9]. We also discuss the relationship between $T = 0$ ensembles

and $T = 0^+$ thermal ensembles. We limit ourselves to systems that are everywhere nonmagnetic.

Interior regions in $\vec{n}$ -space. We first recapitulate Ref. [5] for 1-VR densities, and then generalize to g -VR.

a) $g = 1$. Since there is a non-degenerate non-interacting ground state with a finite gap, corresponding to some $\vec{v}$, we can use non-degenerate perturbation theory to calculate the first-order density change, ${}^1n_l$, due to a weak perturbing potential ${}^1v_l$:

$${}^1n_l = \sum_{l'=1}^{M'} \chi_{ll'}^{-1} {}^1v_{l'}, \quad l = 1, \dots, M'. \quad (1)$$

Since, according to [1], the density $\vec{n}$ of a non-degenerate ground state uniquely determines $\vec{v}$ (no arbitrary constant since $v_M = 0$), the homogeneous set of equations corresponding to (1) has no solution other than ${}^1v_{l'} \equiv 0$, so that (1) can be inverted:

$${}^1v_l = \sum_{l'=1}^{M'} \chi_{ll'}^{-1} {}^1n_{l'} \quad l = 1, \dots, M'. \quad (2)$$

Thus, any first-order change of $\vec{n}$ preserving N produces a density which is also 1-VR. This is the case in the entire M' -D neighborhood surrounding the point $\vec{n}$, in which Δ remains positive.

b) $g > 1$ [10]. We denote by $\Psi_1, \dots, \Psi_g$ a set of g orthogonal degenerate ground-state wave functions corresponding to $\vec{v}^g$, with common energy E and finite gap Δ . The remaining eigenfunctions are $\Psi_{g+1}, \dots$. What conditions must be imposed on infinitesimal $[\mathcal{O}(\lambda)]$ potential changes ${}^1\vec{v}$ which will preserve this degeneracy? In the space of the Ψ_j , the perturbed Hamiltonian matrix is

$$H_{ij} = E_i \delta_{ij} + {}^1V_{ij} \quad i, j = 1, 2, \dots, \quad (3)$$

$E_i = E$ ($i = 1, \dots, g$), $E_i \geq E + \Delta$ ($i \geq g + 1$), and ${}^1V_{ij} \equiv \sum_{l=1}^{M'} \langle i | {}^1v_l | j \rangle$. Because of the finite Δ , the off-diagonal ${}^1V_{ij}$ with $i \leq g$ and $j \geq g + 1$, or vice versa, can be removed to all orders in λ by orthogonal transformations leading to the decoupled $g \times g$ block Hamiltonian

$$\tilde{H}_{ij} = E \delta_{ij} + {}^1V_{ij} + \mathcal{O}(\lambda^2) \quad i, j = 1, \dots, g. \quad (4)$$

Diagonalization by an orthogonal transformation $T_{ij} = \delta_{ij} + {}^1t_{ij} + \mathcal{O}(\lambda^2)$ and the requirement that the eigenvalues remain degenerate gives a transformed Hamiltonian

$$\bar{H}_{ij} \equiv \left(T^{-1} \tilde{H} T \right)_{ij} = (E + {}^1\epsilon) \delta_{ij} + \mathcal{O}(\lambda^2). \quad (5)$$

On inverting the transformation and noting that every matrix commutes with δ_{ij} , one finds $\tilde{H}_{ij} = \bar{H}_{ij} + \mathcal{O}(\lambda^2)$. Comparison with Eq. (4) leads to $\frac{1}{2}(g-1)(g+2)$ conditions of equal diagonal and vanishing off-diagonal ${}^1V_{ij}$. Thus, every point $\vec{v}^g$ with a g -fold degenerate ground state and finite Δ is embedded in a manifold in $\vec{v}$ -space of dimension $D^g = [M' - \frac{1}{2}(g-1)(g+2)]$, in which the degeneracy g and a finite Δ are preserved. Ground-state degeneracies in $\vec{v}$ -space are “rare” in the above sense.

Each $\vec{v}^g$ gives rise to a set of ensemble densities,

$$\vec{n} = \sum_{j=1}^g w_j \vec{n}_j, \quad 0 < w_j \leq 1, \quad \sum_{j=1}^g w_j = 1, \quad (6)$$

with $\vec{n}_j = \vec{n}_j(R^g \Psi_j)$, where R^g is a g -D orthogonal transformation [$\frac{1}{2}g(g-1)$ parameters] and the w_j are normalized weights ($g-1$ parameters). Thus, all $\vec{n}'$ in a finite M' -D neighborhood enclosing a g -VR $\vec{n}$, defined by $\Delta(\vec{n}') > 0$ and $w_j > 0$, are also g -VR: Degeneracy in $\vec{n}$ -space is *not* “rare”.

Boundary surfaces in $\vec{n}$ -space. Except for boundaries of $D \leq M' - 1$, $\vec{n}$ -space is completely filled by M' -D regions S^g , with degenerate ground-state levels and a finite positive gap to the nearest excited state.

Apart from “corners” ($D \leq M' - 2$), interior regions S^g in $\vec{n}$ -space are bounded by $(M' - 1)$ -D internal and external “surfaces” Σ^g , see Fig. 1. Each internal Σ^g separates two interior regions S^g and S^{g+1} . A point $\vec{n}$ on such a Σ^g corresponds to a $(g+1)$ -fold degenerate ground state but with one of the $g+1$ weights equal to zero. On the g -side of Σ^g there is a g -fold degeneracy and a small gap to the $(g+1)$ st state opens up from 0. On the $(g+1)$ -side there is a $(g+1)$ -fold degeneracy and one of the $g+1$ weights starts from 0 and becomes positive.

External boundaries correspond to either one of the n_l becoming zero, reflecting particle conservation; or one of the n_l becoming 1, reflecting the Pauli principle. For examples of both kinds, see the lattice model below.

Imposed symmetry conditions. Consider a Hamiltonian, invariant under a group G of the lattice with irreducible representations h with $D = g_h$. Let the ground state have total degeneracy $g = \sum_h m_h g_h$ due to m_h occurrences of h . If a small perturbation H' respecting the group G is imposed, how many further conditions must be met by H' so that the degeneracy is maintained?

We have shown above that the perturbation must have vanishing off-diagonal matrix elements (MEs) between all degenerate ground states, and equal diagonal MEs.

Off-diagonal MEs. The symmetry of H' assures immediately that all off-diagonal MEs between $h \neq h'$ vanish




FIG. 1: Two interior regions in the M' -D $\vec{n}$ -space, S^g and S^{g+1} , separated by a $(M' - 1)$ -D boundary Σ^g .

[11]. Off-diagonal MEs between two states of the same h involve, by the Wigner-Eckart Theorem, only one independent parameter. The total number of independent parameters characterizing all off-diagonal MEs between the degenerate ground states is therefore $\sum_h m_h(m_h - 1)/2$.

Diagonal MEs. The requirement that all diagonal MEs of H' are equal is automatically satisfied for states belonging to the same h , and involves $\sum_h m_h - 1$ conditions for states belonging to different h .

Thus, to ensure that the g -fold degenerate ground state maintains its full degeneracy requires the imposition of $\sum_h [m_h(m_h - 1)/2 + m_h] - 1$ conditions.

Next we ask: for a ground state with $g = \sum_h m_h g_h$, how many free parameters are there to create ensemble densities invariant under group G of the Hamiltonian?

The most general ensemble density obtainable from g degenerate ground states $\Psi_1, \dots, \Psi_g$ is $\sum_{\mu=1}^g w_{\mu} n(\Psi'_{\mu})$, $\Psi'_{\mu} = R^g \Psi_{\mu}$. However, the Ψ'_{μ} must be chosen in a restricted way to assure that the density has the full symmetry of the Hamiltonian: Let φ_k^{h, ν_h} ($k = 1, \dots, g_h$) be a set of orthogonal eigenfunctions, constituting a basis for a given h of the group G , with occurrence $\nu_h = 1, \dots, m_h$, such that all φ_k^{h, ν_h} transform in the same way under G . The most general density of the required symmetry is

$$\vec{n} = \sum_{h, \nu_h} w^{h, \nu_h} \sum_k \vec{n}(R^{\nu_h} \varphi_k^{h, \nu_h}). \quad (7)$$

The total number of free parameters is given by those in the rotation R^{ν_h} plus the number of independent weights w^{h, ν_h} , i.e. $\sum_h [m_h(m_h - 1)/2 + m_h] - 1$. We observe that the total number of constraints on H' is equal to the total number of free parameters in $\vec{n}$. Therefore the dimensionality of the density invariant under G associated with any set of degenerate states is the same, independent of G .

Lattice Example. We illustrate our topological results in an example including symmetry requirements in $\vec{v}$ - and $\vec{n}$ -space. Consider the singlet ground state of 4 non-interacting electrons, or, equivalently, 2 spinless non-interacting fermions, whose wave functions φ_l^j are defined on a 13-point fcc lattice (see Fig. 2), and vanish on all exterior points. The discrete Schrödinger equation is

$$\left[-\frac{\hbar^2}{2m} \nabla_l^2 + v_l \right] \varphi_l^j = \varepsilon^j \varphi_l^j, \quad j, l = 1, \dots, 13, \quad (8)$$




FIG. 2: 13-point fcc lattice with C_{4h} symmetry: $v_l = V_1$ (V_2) on the sites $l = 1-8$ (9-12), and $v_{13} = 0$.

where, for fcc lattices, $\nabla_l^2 f_l = [\sum_k f_k - 12f_l]/a^2$, with k running over the nearest neighbors of lattice site l . Energies and potentials will be measured in units of $2ma^2/\hbar^2$.

We consider cases of C_{4h} symmetry: $v_l \equiv V_1$, $l = 1-8$; $v_l \equiv V_2$, $l = 9-12$; $v_{13} = 0$. The lattice wave functions belong to the following representations of C_{4h} [11]: A_g (s -like), E_u and A_u (p -like), E_g and B_g (d -like), and B_u (f -like). A_g occurs three times ($1A_g, 2A_g, 3A_g$), E_u occurs twice ($1E_u, 2E_u$), all other representations occur only once. The eigenfunctions and energies required solutions of 3-, 2- and 1-D secular equations.

Consider the ground state of 2 non-interacting spinless fermions. The lowest level is always $1A_g$ ($1s$ -like). The next level depends on V_1 and V_2 , see Fig. 3. The (V_1, V_2) plane divides into three distinct, infinitely extended regions (shown in yellow), in each of which the eigenfunction of the second level belongs to a particular irreducible representation of C_{4h} : Region 1a, $1A_u$ ($2p_z$ -like); region 1b, $2A_g$ ($2s$ -like); and region 1c, $1E_u$ ($2p_x, 2p_y$ -like). *Inside* each region, the ground state is non-degenerate (not counting degeneracies dictated by symmetry, such as in the 2-D representation $1E_u$).

On the boundary lines 2a, 2b and 2c (shown in blue), crossings of levels belonging to different representations of C_{4h} occur, leading to two-fold (accidental) degeneracies. Along 2a, the potential has cubic symmetry ($V_1 = V_2$), and the two lowest p -like levels ($1A_u, 1E_u$) coincide. On 2b, the $2A_g$ and $1A_u$ levels cross, and on 2c, the $2A_g$ and $1E_u$ levels cross. All three boundary lines meet in a single point, “3”, at $V_1 = V_2 = 8$. At this point only, the ground state has a three-fold accidental degeneracy. Note that the degeneracies on 2b and 2c are not due to additional potential symmetries.

As shown above, a ground state with restricted symmetry maintains its degeneracy if $\sum_h [m_h(m_h-1)/2 + m_h] - 1$ conditions are met. In the present example, $m_h = 1$ always. Regions 2a, 2b, 2c are therefore lines (1 condition), region 3 is a point (2 conditions) in the (V_1, V_2) -plane.

We now discuss all $\vec{n}$ with C_{4h} symmetry. Like $\vec{v}$, they are fully characterized by two values, N_1 and N_2 , the total density at points 1-8 and 9-12. $n_{13} = 2 - N_1 - N_2$. N_1 and N_2 are confined within a stripe in the (N_1, N_2) plane, with an “inner” boundary, with $n_{13} = 1$, reflecting




FIG. 3: Top: ground states of 2 non-interacting spinless fermions on the lattice of Fig. 2, with potentials of C_{4h} symmetry. Bottom: associated region of allowed densities.

the Pauli principle ($N_1 + N_2 < 1$ would imply $n_{13} > 1$), and an “outer” boundary, with $n_{13} = 0$, reflecting particle conservation ($N_1 + N_2 \leq 2$).

The manifold of allowed densities in the (N_1, N_2) plane is subdivided into different regions, corresponding to those of the (V_1, V_2) -plane. All of the regions in the (N_1, N_2) plane have $D = 2$. This is in line with our general topological results for g -VR densities.

Fig. 4 illustrates the highest occupied state along paths in the (V_1, V_2) and (N_1, N_2) planes (see dotted lines in Fig. 3 top and bottom): $1A_u$ with weight 1 (region 1a), $1E_u$ with weight 1 (equal-weight combination of $2p_x, 2p_y$ -like states; region 1c), or a linear combination of the degenerate $1A_u$ and $1E_u$ (region 2a), with fractional weights.

Physical Example: the Be-series. Be has a nominal electronic configuration $(1s)^2(2s)^2$ and a term value ^1S . Without interactions, $2s$ and $2p$ are degenerate. The Be-series consists of neutral Be ($Z = 4$) [12], and




FIG. 4: Energies and ensemble weights of the next two levels above $1A_g$, along paths in the (V_1, V_2) and (N_1, N_2) planes (dotted lines in Fig. 3).

the 4-electron ions with $Z = 5, 6, \dots, \infty$. By re-scaling ($\tilde{\mathbf{r}} = Z\mathbf{r}/4$), this is equivalent to a series of Be-like atoms with $Z = 4$ but with weakened interaction strength $\tilde{e}^2 = e^2(4/Z)$. For $Z \rightarrow \infty$, only the nearly degenerate configurations $(1s)^2(2s)^2$ and $(1s)^2(2p)^2$, $(2p)^2 = 3^{-1/2} \sum_{j=1}^3 (2p_j)^2$, need to be considered. This leads to a 2×2 secular equation all of whose MEs are exactly known [13]. The limiting density is explicitly 4-VR:

$$n_0(\tilde{\mathbf{r}}) = 2 \left[\varphi_{1s}(\tilde{\mathbf{r}})^2 + 0.949 \varphi_{2s}(\tilde{\mathbf{r}})^2 + \frac{0.051}{3} \sum_{j=1}^3 \varphi_{2p_j}(\tilde{\mathbf{r}})^2 \right], \quad (9)$$

where the φ 's are scaled hydrogenic wave functions (for $Z = 4$). The configurations $(1s)^2(2s)^2$ and $(1s)^2(2p)^2$ have amplitudes 0.974 and 0.225. (Unlike in CI expansions [14], KS densities consist *strictly* of a *finite* number of orbital densities for *all* Z).

From our previous discussion of boundary surfaces, we observe that the density (9) is *not* on a boundary, since there is no unoccupied degenerate state. Under these conditions, *any* density $n(\mathbf{r})$ sufficiently near $n_0(\mathbf{r})$, in general without any symmetry, is also 4-VR. In particular, there must be a finite range in $1/Z$ of the Be-series where densities are 4-VR. This is borne out by numerical calculations [15].

Density ensembles at $T = 0$ and 0^+ . Let $\vec{v}^g$ be a potential with g degenerate non-interacting, gapped ground states, Φ_j , generating the $\frac{1}{2}(g-1)(g+2)$ -D set of ensemble densities (6). Now introduce a weak per-

turbing potential ${}^1\vec{v}$ of order λ which, in general, splits the degeneracy and leads to eigenstates and eigenvalues $\Psi_j = \bar{R}^g \Phi_j + \mathcal{O}(\lambda)$; $E_j = E + {}^1E_j$, where ${}^1E_j = \mathcal{O}(\lambda)$, and the corresponding densities are $\vec{n}_j$.

For a given low temperature, $\beta^{-1} \ll \Delta$, the canonical density corresponding to $\vec{v} + {}^1\vec{v}$ is given by

$$\vec{n}' = \mathcal{Z}^{-1} \sum_{j=1}^g e^{-\beta {}^1E_j} \vec{n}(\bar{R}^g \Phi_j) + \mathcal{O}(\lambda) + \mathcal{O}(e^{-\beta \Delta}), \quad (10)$$

where $\mathcal{Z} = \sum_j e^{-\beta {}^1E_j}$. This density is equal, to leading order, to the density $\vec{n}$ of Eq. (6), if ${}^1\vec{v}$ is chosen so that $\bar{R}^g \Phi_j = \Psi_j$ and $\mathcal{Z}^{-1} e^{-\beta {}^1E_j} = w_j$. As $\beta \Delta \rightarrow \infty$, ${}^1v \sim \beta^{-1} \rightarrow 0$ and $\vec{n}' \rightarrow \vec{n}$. This establishes the correspondence between ensemble densities at $T = 0$ and canonical densities at low temperatures, $\beta \Delta \gg 1$.

Concluding remarks. The problem of v -representability [3, 4] had cast a shadow of uncertainty over DFT, which was very largely clarified in [7]. Here, building on CCR, we clarify the topology of the $v(\mathbf{r})$ - $n(\mathbf{r})$ mapping. The Be-series and a lattice model are used for illustration. The transition from a dense discrete lattice, r_l , to a continuous variable, $\mathbf{r}$, does not appear to offer difficulties. An open issue is extension to spin magnetism.

Support by NSF Grants No. DMR-96-30452 and DMR-99-76457 and discussions with A. Savin and C. Umrigar are gratefully acknowledged.

* Present Address: Department of Physics, University of Missouri-Rolla, Rolla, MO 65409

- [1] P. Hohenberg and W. Kohn, Phys. Rev. **136 B**, 864 (1964).
- [2] W. Kohn and L. J. Sham, Phys. Rev. **140**, A1133 (1965).
- [3] M. Levy, Phys. Rev. A **26**, 1200 (1982).
- [4] E. H. Lieb, Int. J. Quant. Chem. **24**, 243 (1983).
- [5] W. Kohn, Phys. Rev. Lett. **51**, 1596 (1983).
- [6] Fourier representation is an alternative.
- [7] J. T. Chayes, L. Chayes, and M. B. Ruskai, J. Stat. Phys. **38**, 497 (1985).
- [8] C. A. Ullrich and W. Kohn, Phys. Rev. Lett. **87**, 093001 (2001).
- [9] See also the study of the C_2 molecule by P. R. T. Schipper, O. V. Gritsenko, and E. J. Baerends, Theor. Chem. Acc. **99**, 329 (1998).
- [10] For a basic discussion of the occurrences of degenerate eigenvalues see J. von Neumann and E. Wigner, Phys. Zeitschr. **30**, 467 (1929).
- [11] M. Tinkham, *Group Theory and Quantum Mechanics* (McGraw-Hill, New York, 1964).
- [12] R. van Leeuwen and E. J. Baerends, Phys. Rev. A **49**, 2421 (1994); C. Filippi, C. J. Umrigar, and X. Gonze, Phys. Rev. A **54**, 4810 (1996).
- [13] J. Linderberg and H. Shull, J. Mol. Spectr. **5**, 1 (1960).
- [14] C. F. Bunge, Phys. Rev. **168**, 92 (1968).
- [15] A. Savin, private communication.